

St. John's Jesuit High School & Academy 5901 Airport Highway Toledo, Ohio 43615-7399 419.865.5743 Fax 419.861.5002 www.sjjtitans.org

Date	of	P	lan:	

Diabetes Medical Management Plan

This plan should be completed by the student's personal health care team and parents/guardian. It should be reviewed with relevant school staff and copies s

a place that is easily accessed by the schoauthorized personnel.			-
Effective Dates:			
Student's Name:			
Date of Birth:		etes Diagnosis:	
Grade:		eacher:	
Physical Condition:	Diabetes	s type 2	
Contact Information	·		
Mother/Guardian:			
Address:			
Telephone: Home	Work	Cell	
Father/Guardian:			
Address:			
46			
Telephone: Home	Work	Cell	
Student's Doctor/Health Care Provider:			
Name:			
Address:			
Telephone:	_Emergency Nu	mber:	
Other Emergency Contacts:			
Name:			
Relationship:		5-A-10	
Telephone: Home	Work	Cell	
Notify parents/guardian or emergency co	ntact in the follow	wing situations:	
	,		

Blood Glucose Monitoring
Target range for blood glucose is 70-150 70-180 Other
Usual times to check blood glucose
Times to do extra blood glucose checks (check all that apply)
before exercise
after exercise
when student exhibits symptoms of hyperglycemia
when student exhibits symptoms of hypoglycemia
other (explain):
Can student perform own blood glucose checks?
Exceptions:
Type of blood glucose meter student uses:
Insulin
Usual Lunchtime Dose
Base dose of Humalog/Novolog /Regular insulin at lunch (circle type of rapid-/short-acting insulin used) is units or does flexible dosing using units/ grams carbohydrate.
Use of other insulin at lunch: (circle type of insulin used): intermediate/NPH/lente units or basal/Lantus/Ultralente units.
Insulin Correction Doses
Parental authorization should be obtained before administering a correction dose for high blood
glucose levels. Yes No
units if blood glucose is to mg/dl
units if blood glucose is to mg/dl
units if blood glucose is to mg/dl
units if blood glucose is to mg/dl
units if blood glucose is to mg/dl
Can student give own injections?
Can student determine correct amount of insulin?
Can student draw correct dose of insulin?
Parents are authorized to adjust the insulin dosage under the following circumstances:
read and the second of the sec

For Students with Insulin Pumps Type of pump: Basal rates: 12 am to ____ to ___ _____ to ____ Type of insulin in pump: Type of infusion set: Insulin/carbohydrate ratio: Correction factor: Student Pump Abilities/Skills: Needs Assistance Count carbohydrates Yes No Bolus correct amount for carbohydrates consumed Yes No Calculate and administer corrective bolus Yes No Calculate and set basal profiles Yes No Calculate and set temporary basal rate No Yes Disconnect pump Yes No No Reconnect pump at infusion set Yes Prepare reservoir and tubing Yes No ☐ Yes Insert infusion set No Troubleshoot alarms and malfunctions Yes No For Students Taking Oral Diabetes Medications Type of medication: _____ Timing: _____ Timing: Other medications: Meals and Snacks Eaten at School Is student independent in carbohydrate calculations and management? Yes No Meal/Snack Time Food content/amount Breakfast Mid-morning snack Lunch Mid-afternoon snack Dinner

Yes No

Snack before exercise?

Exercise and Sports A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any:	Foods to avoid, if any: Instructions for when food is provided to the class (e.g., as part of a class party or food sample event): Exercise and Sports A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any: mot exercise if blood glucose level is below mg/dl or if moderate to large urine ketones are present. Hypoglycemia (Low Blood Sugar) Usual symptoms of hypoglycemia: Treatment of hypoglycemia: Glucagon should be given if the student is unconscious, having a seizure (convulsion), or unto swallow. Route , Dosage , site for glucagon injection: arm,thigh, other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are abovemg/dl Treatment for ketones: mg/dl			· · · · · ·		
Instructions for when food is provided to the class (e.g., as part of a class party or food sample event): Exercise and Sports A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any:	Instructions for when food is provided to the class (e.g., as part of a class party or food samp event): Exercise and Sports A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any:	Preferred snack	oods:			
event):	Exercise and Sports A fast-acting carbohydrate such as	Foods to avoid,	f any:			
A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any:	A fast-acting carbohydrate such as should be available at the site of exercise or sports. Restrictions on activity, if any:		_			r food samp
should be available at the site of exercise or sports. Restrictions on activity, if any:	should be available at the site of exercise or sports. Restrictions on activity, if any:	Exercise and S	orts			
mg/dl or if moderate to large urine ketones are present. Hypoglycemia (Low Blood Sugar) Usual symptoms of hypoglycemia: Treatment of hypoglycemia: Glucagon should be given if the student is unconscious, having a seizure (convulsion), or unto swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: mg/dl Treatment for ketones: mg/dl Treatment for ketones: mg/dl	mg/dl or if moderate to large urine ketones are present. Hypoglycemia (Low Blood Sugar) Usual symptoms of hypoglycemia: Treatment of hypoglycemia: Glucagon should be given if the student is unconscious, having a seizure (convulsion), or unto swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	A fast-acting ca should be availa	oohydrate such as le at the site of exercise or spo	orts.		
Usual symptoms of hypoglycemia:	Usual symptoms of hypoglycemia:	Restrictions on not exercise if b	ctivity, if any: ood glucose level is below mg/dl or if moderate	to large urine ketor	mg/dl or at nes are present.	student show
Treatment of hypoglycemia: Glucagon should be given if the student is unconscious, having a seizure (convulsion), or unto swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones: mg/dl	Treatment of hypoglycemia: Glucagon should be given if the student is unconscious, having a seizure (convulsion), or un to swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistance and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Hypoglycemia	Low Blood Sugar)			
Glucagon should be given if the student is unconscious, having a seizure (convulsion), or unto swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones: mg/dl	Clucagon should be given if the student is unconscious, having a seizure (convulsion), or un to swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistance and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Usual sympton				
to swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	to swallow. Route, Dosage, site for glucagon injection:arm,thigh,other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia:	Treatment of hy				
other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	other. If glucagon is required, administer it promptly. Then, call 911 (or other emergency assistant and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:		be given if the student is unco	onscious, having a	seizure (convu	lsion), or un
and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are abovemg/dl Treatment for ketones:	and the parents/guardian. Hyperglycemia (High Blood Sugar) Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Routeother.	Dosage, site for gluca	agon injection:	arm,	thigh,
Usual symptoms of hyperglycemia:	Usual symptoms of hyperglycemia: Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are abovemg/dl Treatment for ketones:			Then, call 911 (or	other emerger	ncy assistanc
Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Treatment of hyperglycemia: Urine should be checked for ketones when blood glucose levels are abovemg/dl Treatment for ketones:	Hyperglycemi	(High Blood Sugar)			
Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Urine should be checked for ketones when blood glucose levels are above mg/dl Treatment for ketones:	Usual symptom				
Treatment for ketones:	Treatment for ketones:	Treatment of hy	erglycemia:			
Treatment for ketones:	Treatment for ketones:	Urine should be	checked for ketones when bloo	od glucose levels a	re above	mg/dl
	Supplies to be Kept at School					
	Supplies to be Kept at School					

Lancet device, lancets, gloves, etc.	
Urine ketone strips	
Insulin pump and supplies	
Insulin pen, pen needles, insulin cartridges	
Fast-acting source of glucose	
Carbohydrate containing snack	
Glucagon emergency kit	
Signatures	
This Diabetes Medical Management Plan has been approv	ed by:
Student's Physician/Health Care Provider	Date
Student's Physician/Health Care Provider	Date
I give permission to the school nurse, trained diabetes personn	
	erform and carry out the diabetes
care tasks as outlined by''s Di Plan. I also consent to the release of the information contained	abetes Medical Management 1 in this Diabetes Medical
Management Plan to all staff members and other adults who h	
who may need to know this information to maintain my child'	•
Acknowledged and received by:	
• • • • • • • • • • • • • • • • • • •	
Student's Parent/Guardian	Date
Student's Parent/Guardian	Date